

Copyright WILEY-VCH Verlag GmbH & Co. KGaA,69469

Weinheim,Germany,2011

NANO MICRO
small

Supporting Information

for *Small*,DOI: 10.1002/sml.201100265

The Passage of Homopolymeric RNA through Small Solid-State Nanopores

Michiel van den Hout , Gary M. Skinner , Sven Klijnhout ,
Vincent Krudde ,and Nynke H. Dekker *

Supplementary Information for:

THE PASSAGE OF HOMOPOLYMERIC RNA THROUGH SMALL SOLID-STATE NANOPORES

Michiel van den Hout, Gary M. Skinner, Sven Klijnhout, Vincent Krudde, and
Nynke H. Dekker[§]

[§]Kavli Institute of Nanoscience Delft
Delft University of Technology
Lorentzweg 1
2628 CJ Delft
The Netherlands
e-mail: N.H.Dekker@tudelft.nl
Tel: +31-(0)15-27 83219
Fax: +31-(0)15-27 81202




Figure S1. Dependence of the average dwell times of the H- (closed circles) and L-events (open circles) on nanopore diameter. The average dwell time for both event types gradually increases with decreasing nanopore diameter. The variation in the data is large, likely resulting from the large variation in length of the poly(U) molecules (see Molecule Synthesis, in the Experimental Section of the main text).